



LabCorp Dublin  
6370 Wilcox Road  
Dublin, OH 43016-1269

Phone: 800-282-7300

Specimen Number <b>354-127-9542-0</b>		Patient ID		Control Number	Account Number 90000045	Account Phone Number 336-436-8645	Route 00
<b>SAMPLE REPORT</b>				Account Address			
Patient Last Name				LabCorp Test Master			
Patient First Name <b>500183</b>		Patient Middle Name		Test Account			
Patient SS#		Patient Phone		3060 South Church Street			
Age (Y/M/D)		Date of Birth		Sex N		Fasting	
Patient Address				Additional Information			
				NORMAL			
Date and Time Collected 12/19/16 00:00	Date Entered 12/19/16	Date and Time Reported		Physician Name	NPI	Physician ID	

Tests Ordered							
Anti-Mullerian Hormone (AMH)							

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
-------	--------	------	-------	--------------------	-----

<b>Anti-Mullerian Hormone (AMH)</b>					01
Anti-Mullerian Hormone (AMH)	9.50		ng/mL		
For assays employing antibodies, the possibility exists for interference by heterophile antibodies in the samples. <sup>1</sup>					
1. Kricka L. Interferences in Immunoassays - still a threat. Clin. Chem. 2000; 46: 1037-1038.					
Reference Range:					
Females 20 - 25y: 1.23 - 11.51					
Median 4.70					
AMH concentrations of $\geq 1.06$ ng/mL is correlated with a better response to ovarian stimulation, produced more retrievable oocytes and higher odds of live birth according to Gleicher et al. Fertility and Sterility. 2010: 94:2824-2827. The current AMH test method correlates with the study method with a slope of 0.94.					
Females at risk of ovarian hyperstimulation syndrome or polycystic ovarian syndrome (PCOS) may exhibit elevated serum AMH concentrations. AMH levels from PCOS patients may be 2 to 5 fold higher than age-appropriate reference interval values.					
Granulosa cell tumors of the ovary may secrete AMH along with other tumor markers. Elevated AMH is not specific for malignancy, and the assay should not be used exclusively to diagnose or exclude an AMH-secreting ovarian tumor.					

01	ES	Esoterix Endocrinology	Dir: Samuel Pepkowitz, MD
		4301 Lost Hills Road, Calabasas Hills, CA 91301-5358	
For inquiries, the physician may contact <b>Branch: 800-222-7566 Lab: 800-282-7300</b>			

<b>SAMPLE REPORT, 500183</b>		<b>354-127-9542-0</b>	Seq # 0000
------------------------------	--	-----------------------	------------



LabCorp Dublin  
6370 Wilcox Road  
Dublin, OH 43016-1269

Phone: 800-282-7300

Specimen Number <b>354-127-9543-0</b>		Patient ID		Control Number	Account Number 90000045	Account Phone Number 336-436-8645	Route 00
<b>SAMPLE REPORT</b>				Account Address			
Patient Last Name				LabCorp Test Master			
Patient First Name <b>500183</b>		Patient Middle Name		Test Account			
Patient SS#		Patient Phone		3060 South Church Street			
Age (Y/M/D)		Date of Birth		Sex N		Fasting	
Patient Address				Additional Information			
				ABNORMAL			
Date and Time Collected 12/19/16 00:00		Date Entered 12/19/16		Date and Time Reported		Physician Name	
						NPI	
						Physician ID	

Tests Ordered							
Anti-Mullerian Hormone (AMH)							

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
-------	--------	------	-------	--------------------	-----

<b>Anti-Mullerian Hormone (AMH)</b>					
Anti-Mullerian Hormone (AMH)	>15.0		ng/mL		01
<p>For assays employing antibodies, the possibility exists for interference by heterophile antibodies in the samples.<sup>1</sup></p> <p>1. Kricka L. Interferences in Immunoassays - still a threat. Clin. Chem. 2000; 46: 1037-1038.</p> <p>Reference Range:</p> <p>Females 20 - 25y: 1.23 - 11.51</p> <p>Median 4.70</p> <p>AMH concentrations of <math>\geq 1.06</math> ng/mL is correlated with a better response to ovarian stimulation, produced more retrievable oocytes and higher odds of live birth according to Gleicher et al. Fertility and Sterility. 2010: 94:2824-2827. The current AMH test method correlates with the study method with a slope of 0.94.</p> <p>Females at risk of ovarian hyperstimulation syndrome or polycystic ovarian syndrome (PCOS) may exhibit elevated serum AMH concentrations. AMH levels from PCOS patients may be 2 to 5 fold higher than age-appropriate reference interval values.</p> <p>Granulosa cell tumors of the ovary may secrete AMH along with other tumor markers. Elevated AMH is not specific for malignancy, and the assay should not be used exclusively to diagnose or exclude an AMH-secreting ovarian tumor.</p>					

01	ES	Esoterix Endocrinology	Dir: Samuel Pepkowitz, MD
		4301 Lost Hills Road, Calabasas Hills, CA 91301-5358	
For inquiries, the physician may contact <b>Branch: 800-222-7566 Lab: 800-282-7300</b>			

<b>SAMPLE REPORT, 500183</b>		<b>354-127-9543-0</b>	Seq # 0000
------------------------------	--	-----------------------	------------