

Specimen ID:

Acct #:

Phone:

Rte:

Control ID:

Patient Details

DOB:
Age(y/m/d):
Gender: SSN:
Patient ID:

Specimen Details

Date collected: 05/10/2018 1300 Local
Date received: 05/10/2018
Date entered: 05/10/2018
Date reported: 05/11/2018 1210 ET

Physician Details

Ordering:
Referring:
ID:
NPI:

General Comments & Additional Information

Alternate Control Number:
Total Volume: Not Provided
Ordered Items

Alternate Patient ID:
Fasting:

CEA; Prostate-Specific Ag, Venipuncture

LAB	TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	
CEA		2.4		ng/mL	0.0 - 4.7	01
		Roche ECLIA methodology			Nonsmokers <3.9 Smokers <5.6	

Prostate-Specific Ag, Serum

Prostate Specific Ag, Serum 0.8 ng/mL 0.0 - 4.0 01

Roche ECLIA methodology.

According to the American Urological Association, Serum PSA should decrease and remain at undetectable levels after radical prostatectomy. The AUA defines biochemical recurrence as an initial PSA value 0.2 ng/mL or greater followed by a subsequent confirmatory PSA value 0.2 ng/mL or greater.

Values obtained with different assay methods or kits cannot be used interchangeably. Results cannot be interpreted as absolute evidence of the presence or absence of malignant disease.

01 PDLCA LabCorp Phoenix
5005 S 40th Street Ste 1200, Phoenix, AZ 85040-2969

Dir: Earle Collum, MD

For inquiries, the physician may contact **Branch: 800-762-4344 Lab: 800-788-9743**